

NDVI-based productivity and heterogeneity as indicators of plant-species richness in boreal landscapes

Miia Parviainen^{1)*}, Miska Luoto¹ and Risto K. Heikkinen²

¹⁾ Department of Geography, P.O. Box 3000, FI-90014 University of Oulu, Finland (*e-mail: miia.parviainen@oulu.fi)

²⁾ Finnish Environment Institute, Research Programme for Biodiversity, P.O. Box 140, FI-00251 Helsinki, Finland

Received 22 Dec. 2008, accepted 6 May 2009 (Editor in charge of this article: Jaana Bäck)

Parviainen, M., Luoto, M. & Heikkinen, R. K. 2010: NDVI-based productivity and heterogeneity as indicators of plant-species richness in boreal landscapes. *Boreal Env. Res.* 15: 301–318.

Decision making in managing biological diversity is critically dependent on adequate information concerning species-richness patterns and a rigorous understanding of species–habitat relationships. Measures of primary productivity derived from satellite images may provide useful cost-effective estimates of species richness and distribution patterns over wide areas. We constructed Generalized Additive models (GAM) to investigate the potential of primary productivity and its heterogeneity based on Normalized Difference Vegetation Index (NDVI) to explain the species richness in 28 separate vascular plant families in boreal forest landscapes, northern Finland. The productivity models explained on average more of the species richness than the heterogeneity models. However, models that performed best were produced by combining productivity and heterogeneity variables into the same models. Species richness responded mainly unimodally or positively to productivity and its heterogeneity. We conclude that measures of productivity and heterogeneity based on remote sensing can provide useful ‘first filters’ of locations of high diversity in plant families in boreal landscapes.

Introduction

Developing means for rapid forecasting of species richness and distributions using a few easily measured environmental variables is increasingly important in the assessment of impacts of anthropogenic and natural disturbances on biodiversity under limited resources (Kerr and Ostrovsky 2003, Seto *et al.* 2004). Remote sensing offers an inexpensive means to derive spatially complete environmental information for large areas in a consistent and regular manner (Muldavin *et al.* 2001, Foody and Cutler 2003).

Such information may provide valuable tools for the prediction of spatial patterns of biodiversity attributes.

The current paradigm is that climate governs species distribution and richness patterns on broad biogeographical scales (Currie 1991, Huntley *et al.* 1995, Parmesan 1996, H-Acevedo and Currie 2003, Thuiller *et al.* 2004), whereas land cover and spatial distribution of suitable biotopes determine species occupancy patterns more than climate at finer spatial resolutions (Dunning *et al.* 1992, Opdam and Wascher 2004, Pearson *et al.* 2004, Thuiller *et al.* 2004). The

amount of energy available in a system, often measured as primary productivity, is also considered as one of the major determinants of biodiversity, especially species richness (Currie 1991, Rosenzweig 1995, Hawkins and Porter 2003). The Normalized Difference Vegetation Index (NDVI hereafter) (Tucker 1979) is one of the most extensively used remote-sensing based vegetation indices (for more discussion *see* Gould 2000, Nagendra 2001, Kerr and Ostrovsky 2003, Pettorelli *et al.* 2005). Remotely sensed NDVI observations provide direct estimates of primary productivity because NDVI measures the energy entering the ecosystem (Tucker and Sellers 1986, Reed *et al.* 1994, Paruelo *et al.* 1997, Mittelbach *et al.* 2001, Benayas and Scheiner 2002, Levin *et al.* 2007). Due to this intimate link between NDVI and primary productivity, NDVI has been found to be a useful predictor of regional variation in species richness (e.g. Gould 2000). NDVI is related to certain critical environmental factors (Seto *et al.* 2004), which improves its potentiality as a biodiversity predictor.

The relationship between primary productivity and species richness is generally assumed to be hump-shaped (Grime 1973, Rosenzweig and Abramsky 1993, Waide *et al.* 1999, Mittelbach *et al.* 2001, Fairbanks and McGwire 2004), but other response shapes have also been observed (Mittelbach *et al.* 2001), including positive (Rosenzweig and Abramsky 1993, Rosenzweig 1995, Mittelbach *et al.* 2001, Whittaker and Heegaard 2003, Gillespie 2005) and negative ones (Oindo and Skidmore 2002). The position of the optimum productivity for species richness may be different for different species, ecosystems, observation scales and biogeographical regions (Waide *et al.* 1999).

Apart from the maximum or intermediate productivity, species richness also responds to variation in primary productivity (Rosenzweig 1995, Kerr and Packer 1997, Hawkins and Porter 2003, Bailey *et al.* 2004). Heterogeneity in environmental factors is known to contribute to the diversity of plant communities (Ricklefs 1977, Grime 1979, Palmer 1994, Grace 1999, Lundholm and Larson 2003). Increase in heterogeneity implies an increase in niches, allowing more species to coexist. Spectral heterogeneity

of remotely sensed images is related to the spatial heterogeneity of the environment, in particular to the plant species richness (Rocchini *et al.* 2004). For example, Gould (2000) showed that variation in NDVI can help in explaining the variation in species richness in arctic landscapes of Canada. The studies by Palmer *et al.* (2002), Rocchini *et al.* (2004), Rocchini *et al.* (2005) and Rocchini (2007) also revealed a significant relationship between heterogeneity and species richness, in their case from Mediterranean landscapes. The relationship between variation of NDVI in space and/or time (“heterogeneity in NDVI”) and species richness has generally been found to be positive (Gould 2000, Oindo and Skidmore 2002, Fairbanks and McGwire 2004, Levin *et al.* 2007).

Several studies have highlighted the relevance of remote sensing information in biodiversity modeling (Gould 2000, Nagendra 2001, Kerr and Ostrovsky 2003, Pettorelli *et al.* 2005). However, there has been little effort to use such remote-sensing based estimators of habitat heterogeneity to predict species richness at the mesoscale, i.e. spatial resolutions ranging from 0.5 km to 2 km. Moreover, studies comparing predictions of species richness based on heterogeneity in NDVI with those based on NDVI primary-productivity values are largely lacking. In this study, we used mean and maximum NDVI values as a measure of primary productivity, and the range and standard deviation of NDVI values as a measure of the heterogeneity of productivity. The specific aims were to investigate the potential of (1) primary productivity and (2) heterogeneity to explain the richness patterns in 28 vascular plant families in a high-latitude forest landscape, northern Finland using plant species richness data from a spatial grid system (440 squares of 25 ha within an area of 110 km²). We modeled the species richness in 28 plant families separately instead of using overall total species richness. This was because we were interested in investigating whether certain plant families are more closely related to NDVI-based predictors than others, and because different plant families may exhibit contradicting responses to productivity, making the overall richness–productivity relationships somewhat blurred.




Fig. 1. A map of the richness of plant families in each of the 25-ha grid squares in the study area. The total number of families is 28.

Material and methods

Study area

The study area consists of 440 grid squares 25 ha in size and located in Oulanka National Park in northern Finland (66°22'N, 29°19'E) (Fig. 1). Oulanka National Park (total area = 27 746 ha) was founded in 1956. Oulanka National Park is located near the southern edge of the northern boreal-forest zone (Parviainen *et al.* 2008). The northern part of the study area is characterised by large open mires, whilst the southern part is dominated by forested hills, but also with mosaic of river valleys, water bodies, open bogs and mires. Dominant tree species are Norway spruce (*Picea abies*) and Scots pine (*Pinus sylvestris*) and birches (*Betula* spp.). The vegetation is relatively rich with arctic, eastern, Siberian and southern species (Vasari *et al.* 1996). The mean annual temperature is ca. 0.5 °C, the growing season lasts ca. 128 days and the difference

between the mean temperature of the coldest (January average ca. -14 °C) and the warmest (July average ca. 15 °C) month is ca. 29 °C (Atlas of Finland 1987). The climate in the region is thus more continental than in most other parts of northern Europe but with a maritime (humid) element added. Topography varies conspicuously and elevation ranges from 140 to 440 m a.s.l.

Plant family data

This study used the records of plant species found in the flora survey made by Söyrinki and Saari in the year 1980 (Söyrinki and Saari 1980). A total of 476 plant species were recorded in the survey. Plant data included detailed information on the geographical location of the occurrences (coordinates in the uniform grid system, Grid 27°E).

The data on the plant family richness (taxonomy follows Hämet-Ahti *et al.* 1986) for the 440 25-ha grid cells constituting the study area

were generated by calculating the number of species representing a certain plant family separately for each of the 28 plant families from the original georeferenced data of Söyrinki and Saari (1980). The plant family data for the study area consisted of richness records of 69 plant families. Of these families, 28 families with four or more records in the 440 grid squares of 25 ha and covering the whole study area were used in the analyses (Fig. 1 and Table 1). In total, 359 vascular plant species were included in the study. The most species-rich plant families were the Cyperaceae (58 species), Poaceae (47 species) and Asteraceae (22 species) (*see* Appendix 1 for a complete species list).

We are aware of the fact that there is a slight temporal mismatch between the plant species data (year 1980) and Landsat-imagery (year

2000). However, we assume that the effect of the temporal mismatch is only minor, since study area is a national park, where vegetation changes are relatively slow and human impacts very unsubstantial. Furthermore, the plant families found in the 1980s have principally been found also in later reinventories made in the same area.

Remote sensing data

The Normalized Difference Vegetation Index (NDVI) (Rouse *et al.* 1973, Tucker 1979) is the most frequently used parameter for quantifying productivity and above-ground biomass of ecosystems (Tucker 1979, Box *et al.* 1989). It is based on the strong absorption of incident radiation by chlorophyll in the red (RED) spectral region, and the contrasting high reflectance by plant cells in the near infrared (NIR) region. Because NDVI is based on the normalized ratio of the reflectance in these two spectral regions (*see* the formula below), it is an indicator of the greenness of vegetation canopies, which enables separation of vegetation from other land coverage.

The NDVI was calculated for each 25-ha grid square using the formula:

$$\text{NDVI} = (\text{NIR} - \text{RED}) / (\text{NIR} + \text{RED})$$

NDVI measures were generated from one geo-corrected Landsat ETM+ satellite scene (Table 2 and Fig. 2). The scene was acquired in July 2000 to coincide with the growing season. First, the Landsat scene was rectified according to digitized topographic maps (scale 1:20 000). The geometric correction of the planimetric root-mean-square error (RMSE) of test ground control points of the image was 7.9 m. The spatial resolution of the rectified Landsat TM image was set as 25 m, and new values for the pixels were resampled using a cubic convolution interpolation method (Hjort and Luoto 2006). Second, the image was topographically corrected using the 'Ekstrand correction method' (Ekstrand 1996). Finally, in order to decrease the effects of atmospheric variations, the image was atmospherically corrected using the SMAC algorithm, which is a semi-empirical correction method developed at the VTT, Technical Research Center of Finland

Table 1. The total, minimum, mean and maximum numbers of species in the 28 vascular plant families, recorded from the 440 25-ha grid squares situated in Oulanka National Park, northern Finland.

Family	Total	Min	Mean	Max
Lycopodiaceae	5	0	1.9	4
Equisetaceae	8	0	4.1	8
Dryopteridaceae	13	0	1.7	8
Potamogetonaceae	9	0	0.6	5
Juncaceae	10	0	1.9	6
Poaceae	47	0	10.4	36
Cyperaceae	58	0	14.7	29
Orchidaceae	13	0	2.1	10
Salicaceae	16	0	5.2	10
Betulaceae	4	1	3.3	4
Polygonaceae	9	0	0.7	6
Caryophyllaceae	19	0	0.8	8
Ranunculaceae	14	0	2.8	10
Brassicaceae	12	0	0.5	8
Saxifragaceae	4	0	0.2	4
Rosaceae	16	0	6.8	13
Fabaceae	9	0	0.6	7
Violaceae	5	0	1.3	5
Onagraceae	6	0	1.5	5
Apiaceae	6	0	1.0	4
Pyrolaceae	6	0	2.6	6
Ericaceae	10	4	7.3	9
Rubiaceae	5	0	1.4	4
Lamiaceae	5	0	0.4	4
Scrophulariaceae	11	0	3.5	8
Lentibulariaceae	6	0	1.1	5
Asteraceae	22	0	3.4	13
Cichoriaceae	11	0	1.8	7
Total	359			




Fig. 2. Environmental variables calculated for each of the 25-ha grid squares: (A) mean NDVI, (B) maximum NDVI, (C) standard deviation of NDVI and (D) range of NDVI.

(Hjort and Luoto 2006). The satellite image was provided by the Finnish Environment Institute (SYKE) and orthorectified by METRIA, Sweden (Härmä *et al.* 2004).

Statistical analyses

GAMs

The 28 separate response variables, i.e. the number of species in each of the 28 vascu-

lar plant families, were related to the explanatory variables with generalized additive models (GAMs) (Hastie and Tibshirani 1990) using the GRASP ver. 3.2 package (Lehmann *et al.* 2002) for S-Plus ver. 6.1 (Insightful Corp., Seattle, WA, USA). GAMs are nonparametric extensions of generalized linear models (McCullagh and Nelder 1989). They have extensively been used in ecological applications (*see* Yee and Mitchell 1991, Guisan *et al.* 2002), because they permit both parametric and non-parametric additive response shapes, as well as a combination of the

Table 2. List of remote sensing variables (Landsat ETM+) used as explanatory variables in the modeling.

	Remote sensing variable	Abbreviation	Min	Max	Mean	SD
Productivity	Normalized Difference Vegetation Index (mean)	NDVI mean	0.067	0.656	0.442	0.101
	Normalized Difference Vegetation Index (max)	NDVI max	0.404	0.935	0.769	0.093
Heterogeneity	Normalized Difference Vegetation Index (SD)	NDVI SD	0.061	0.486	0.131	0.052
	Normalized Difference Vegetation Index (range)	NDVI range	0.354	1.723	0.752	0.275




Fig. 3. (A) The amount of explained deviance (%) and (B) the cross-validated correlation values in the studied 28 plant families, derived from three different modeling settings. The boxes show medians, and 1st and 3rd quartiles values.

two within the same model (Wood and Augustin 2002). The GAM models were built using a stepwise selection procedure to select relevant explanatory variables, starting with a full model in which all predictors are fitted and alternately omitting and re-introducing one model component at each step. A Poisson probability distribution was selected for the response (count data), and the link function was set to logit. A smoothing spline method was chosen to smooth the studied variables, using 1 or 4 degrees of freedom (df) by default (Venables and Ripley 2002).

GAMs were fitted using three sets of explanatory variables for each of the 28 plant family richnesses. The first model for each family was built using mean and maximum values of NDVI in 25 ha grid squares; hereafter the productivity model. The second model was based on standard deviation and range in the NDVI values; hereafter the heterogeneity model. The final model (hybrid model) included all the productivity and heterogeneity variables.

The performance of models was evaluated using (1) the percentage of explained deviance (D^2), and (2) the correlation coefficient (r) between the observed and predicted values based on a four-fold cross-validation. The folds were generated by dividing the entire data set into four subsets, so that each subset contained an equal number of randomly selected data points. Each subset was dropped in turn from the data set, the

model was calibrated with the three other subsets and the predictions were made for the omitted data points. After recombining the four folds, predictions were plotted against observed data (Lehmann *et al.* 2002).

Additionally, we validated visually the response curves to identify the correlation structure between response and remote sensing variables. Wilcoxon signed-rank tests were used to compare the performance of different competing models.

Results

The productivity models explained on average 19.5% of plant species richness and the heterogeneity models 12.9% (Fig. 3 and Appendix 2). The Wilcoxon signed-rank test revealed significant differences between these two models both in explained deviance (Wilcoxon signed rank test: $z = -3.652$, $p < 0.001$) and cross-validated correlation values ($z = -3.268$, $p = 0.001$) in favor of the productivity models. The productivity model explained best the richness of the Brassicaceae family (40.0%) and the heterogeneity model that of the Potamogetonaceae family (30.6%) (Appendix 2). Heterogeneity models for the families Equisetaceae (9.90%), Cyperaceae (8.95%) and Lentibulariaceae (7.51%) accounted for a higher amount of explained deviance than

the corresponding productivity models (7.09%, 2.72% and 1.97%, respectively), and the same also applied for cross-validated correlations (Appendix 2). Cross-validation correlation ratios of productivity and heterogeneity models were on average moderate, with mean r values of 0.40 with the productivity models and 0.32 with the heterogeneity models (Fig. 3 and Appendix 2).

Incorporating heterogeneity variables alongside productivity variables into the models improved the mean explanatory power (18.2%) and cross-validated correlation values (12.0%) statistically significantly (Wilcoxon signed rank test: $p < 0.001$) (Fig. 3 and Appendix 2). The highest improvement both in explained deviance and cross-validated correlation ratios was obtained with the families Cyperaceae (over 400% improvement in explained deviance and over 800% improvement in cross-validated correlation ratios compared to the productivity model) and Lentibulariaceae (over 300% and 200% improvements, respectively) (Appendix 2). Models for the families of Brassicaceae (42.1%), Lamiaceae (41.1%), and Potamogetonaceae (36.2%) gave on average the highest explained deviance ratios. The highest cross-validated correlations were obtained with the families Ranunculaceae ($r = 0.644$) and Lamiaceae ($r = 0.603$) (Appendix 2).

The range of NDVI values was the most frequently selected predictor of plant species richness. It was selected in 93% of the hybrid models (26 out of 28), whereas the mean and maximum values of NDVI both entered into 82.1% (23 out of 28) of the models (Table 3). A humped relationship between mean productivity (mean NDVI) and species richness was found in 60.9% of these cases (14 out of 23), followed by positive relationships (9 out of 23, 26.1%), as e.g. in the case of Orchidaceae (Fig. 4A) and Lycopodiaceae (Fig. 4B). In total, 20 models out of 23 (87.0%) found a positive relationship between maximum productivity measured as maximum NDVI and species richness, as compared with 2 hump-shaped (8.7%) and one negative (4.3%) relationships. The relationships between species richness and standard deviation in NDVI were mostly hump-shaped (8 out of 14, 57.1%). However, positive and hump-shaped relationships were equally common (12 out of

26, 46.2%) when heterogeneity was measured as range in NDVI values (Table 3). Species richness was associated negatively in 35.7% (5 out of 14) of the cases with standard deviation of NDVI, and in 7.7% (2 out of 26) of the cases with the range of NDVI. As an example, the shapes of the responses in univariate models for the families Orchidaceae and Lycopodiaceae are presented in Fig. 4.

In general, the highest Spearman rank correlations are between NDVI max and richness values, whereas the correlations between NDVI mean and richness values are typically rather low (*see* Table 4).

Table 3. Summary of the response shapes between the 28 plant species families and each remote sensing variable in the hybrid GAM models. For abbreviations of the remote sensing variables *see* Table 2. The direction of the effect is indicated with symbols (+ = positive linear correlate; - = negative linear correlate; \cap = non-linear correlate with a hump-shaped response curve).

Family	NDVI			
	Mean	Max	SD	Range
Lycopodiaceae	+	+		
Equisetaceae	+			\cap
Dryopteridaceae	+	+	-	+
Ranunculaceae	\cap	+	\cap	+
Betulaceae	\cap			
Caryophyllaceae	\cap	+	\cap	+
Polygonaceae		+		\cap
Violaceae	\cap	+		+
Brassicaceae	\cap	+		+
Salicaceae	\cap		\cap	\cap
Ericaceae		\cap		\cap
Pyrolaceae	+	+	-	\cap
Saxifragaceae	\cap	+		\cap
Rosaceae	\cap	+	\cap	\cap
Fabaceae	\cap	+	\cap	+
Onagraceae	+		-	-
Apiaceae		+		+
Rubiaceae	\cap	+	+	+
Lamiaceae	\cap	+		+
Scrophulariaceae	\cap	+		+
Lentibulariaceae		-		\cap
Asteraceae	+	+		\cap
Cichoriaceae	+	+	-	\cap
Orchidaceae	+		-	-
Potamogetonaceae	\cap	+	\cap	+
Juncaceae		+		\cap
Cyperaceae	\cap	\cap	\cap	\cap
Poaceae	+	+	\cap	+
Count	23	23	14	26




Fig. 4. Response shapes in the univariate GAM models for the families of **(A)** Orchidaceae and **(B)** Lycopodiaceae. The dashed lines are approximate 95% point-wise confidence intervals; tickmarks show the location of observations along the variable range; y-axis represents the effect of the respective variable; s represents the smooth term of GAM. For abbreviations of the environmental variables see Table 1.

Discussion

Vascular plant species richness is an important ecosystem feature and one attribute that characterizes the biodiversity of an area (Currie 1991, Gaston 2000, Gould 2000). The high-latitude environment restricts the presence and productivity of vascular plant species. In particular, low temperature, short growing season, low rates of soil nutrient cycling and extremes of soil moisture may limit primary productivity (Bonan and Shugart 1989). Therefore high-latitude landscapes are often considered to be relatively

homogeneous in the structure of their vegetation and rather poor in species richness. However, such landscapes may comprise very different plant assemblages (Virtanen *et al.* 2006). For example, herb-rich forests within the boreal zone may be considerably rich in terms of their species diversity (Kuusipalo 1984, Heikkinen 1998), although they may cover only a fraction of the total forest area (Airaksinen and Karttunen 2001). Such areas are particularly important for the maintenance of biological diversity in boreal landscapes. Mesoscale biodiversity patterns are an important component of the diversity that occurs in a landscape or region, and represent also the scale at which management decisions are often made in terms of land use and conservation (Hill and Keddy 1992, Stoms 1994, Heikkinen 1998, Gould 2000).

Table 4. The Spearman rank correlation between NDVI based productivity and heterogeneity indices and richness of all plant species, herbaceous species, tree species and family richness (number of different plant families) in the studied 440 25-ha grids.

Richness	NDVI			
	Mean	Max	SD	Range
All plant species	0.125**	0.456***	0.188***	0.396***
Tree species	0.218***	0.413***	0.031 ^{NS}	0.252***
Herbaceous species	0.075 ^{NS}	0.417***	0.225***	0.414***
Family richness	0.159**	0.498***	0.167***	0.376***

*** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$; ^{NS} = non significant.

Explanatory power of the models

Our results suggest that NDVI provides a useful synthetic predictor that is significantly related to environmental variables affecting the richness of vascular plant species. Especially productivity measured as the mean and maximum NDVI values appears to account for a considerable amount of species richness within the different plant families. This is in agreement with earlier findings indicating that primary productivity is

one of the major determinants of species richness (Currie 1991, Rosenweig 1995, Hawkins and Porter 2003). The same authors have also showed that heterogeneity also often correlates positively with species richness. Heterogeneity of NDVI appears indeed to have a high predictive power for the species richness of a given site, across a range of spatial scales (Gould 2000, Kerr *et al.* 2001, Oindo and Skidmore 2002, Rocchini *et al.* 2004). Areas with a wide range of NDVI values, or with high standard deviations of the mean NDVI, are places where dense vegetation cover alternates with sparsely vegetated sites, or environments with different successional stages of vegetation — both of which may indicate high heterogeneity. However, in our study heterogeneity models explained on average less of the plant family species richness than the productivity models. This suggests that productivity is a stronger descriptor of mesoscale plant family richness patterns in high-latitude forest landscapes than heterogeneity (*see also* Levin *et al.* 2007). This result may be related to the scale-dependence of the relationship between heterogeneity and species richness (Palmer *et al.* 2002, Rocchini *et al.* 2004). As the size of sample plots increases, the importance of within-plot heterogeneity increases (Grace 1999). For example, Rahbek and Graves (2001) suggested that habitat heterogeneity may be a more significant determinant of larger scale diversity in some regions. However, our study area is located in boreal high latitude landscapes which are a relatively harsh environment for the species as compared with more southern temperate and tropical environments. A recent review by Hawkins *et al.* (2003) indicated that, at least on broad biogeographical spatial scales, energy- and productivity-related factors often play the most critical role in determining species richness of different organism groups especially in northern landscapes in Canada, Siberia and northern Europe. This is because available energy acts as the most critical limitation and thus constitutes the primary constraint for species richness in such environments.

However, incorporating heterogeneity variables alongside productivity variables improved the plant species richness models. Although the absolute increases in the amount of explained deviance and cross-validated correlations were

generally small, they showed a clear trend and were statistically significant. Moreover, although the relative contribution of heterogeneity variables in our modeling experiment was rather low, they were selected into the majority of the models; 26 out of 28 models included at least one heterogeneity variable. Thus although productivity appears to be the prominent factor in explaining richness patterns, richness in most of the plant families can be increased due to increased habitat (productivity) heterogeneity. Physically and biologically more diverse environments enable finer-scale utilization of limited resources by different species in relatively small areas (Hawkins *et al.* 2003).

In summary, our results indicate that both productivity and heterogeneity play an important role in determining mesoscale species richness in high-latitude forest landscapes, because they reflect different aspects of productivity. In a similar vein, Foody and Cutler (2003) pointed out that more accurate information could be obtained using more complex approaches, e.g. by using standard deviation of NDVI alongside the mean values. The importance of productivity vs. heterogeneity appears to vary between plant families differing in their habitat requirements. For example, the explanatory capacity of the heterogeneity models was substantially higher than that of the productivity models in the case of the families Equisetaceae, Potamogetonaceae and Lentibulariaceae. These families are known to prefer aquatic and shore habitats, where resource heterogeneity may be rather high.

The unexplained variation in our richness models was relatively high, which can be at least partly arise from unmeasured environmental variables (Austin 2002). In addition to productivity, other factors also affect species richness. One of these factors is topography (elevation, slope, aspect) (Austin 2002, Virtanen *et al.* 2006). Relationships between richness patterns and various ecological, geographical or other factors have been dealt in many papers (e.g. Currie 1991, Borcard *et al.* 1992, Heikkinen 1996, Iverson *et al.* 1997, Luoto 2000, Luoto *et al.* 2002, Wiens and Donoghue 2004). Thus, in detailed analyses of plant species richness in a landscape, several additional factors other than merely productivity should be taken into account.

Response shapes

Productivity measured as the mean NDVI values played a significant role in explaining species richness, showing mainly a hump-shaped species–energy relationship, where richness is reduced by abiotic stress at one end of the productivity gradient and by competition at the other (Grime 1979). Our findings are in line with those of former studies, in which a hump-shaped relationships between primary productivity and species richness have often been observed, especially at regional and smaller scales (*see* Currie and Paquin 1987, Wright *et al.* 1993 for more details, Rosenweig 1995, Grace 1999, Waide *et al.* 1999, Gaston 2000, Grime 2001, Mittelbach *et al.* 2001). Grace (1999) reviewed the literature on productivity–diversity relationships in herbaceous plant communities and concluded that most studies relating plant species richness to plant biomass in small plots showed a hump-shaped relationship. Mittelbach *et al.* (2001) also found that a unimodal relationship was detected most frequently when patterns were analyzed across vascular plant community types. At the individual species level, Waide *et al.* (1999) found that whereas hump-shaped relationships were common for some taxa and at some geographical and ecological scales, other patterns also occurred. Competition is probably the most reasonable explanation for a decline in species richness at high productivity levels, but other causes have also been proposed, including facilitation (Michalet *et al.* 2006), biogeographical affinity (Harrison and Grace 2007), differences in dispersal probability (Pärtel *et al.* 2007) and varying evolutionary processes (VanderMeulen *et al.* 2001, Bruun and Ejrnæs 2006). Furthermore, one potential explanation for hump-shaped patterns may be that spatial heterogeneity in resource availability is maximal at intermediate productivity, leading to the prevention of competitive exclusion between plant species (Tilman 1982). However, plant family richness was not always unimodally related with productivity. Many of the plant families, for example Dryopteridaceae, Orchidaceae, Ranunculaceae and Pyrolaceae, represented monotonically linear, positive association with the mean NDVI.

It should be noted that mean productivity reflects the average amount of biomass or resource variables, whereas maximum NDVI is more related to the highest potential productivity. In stable, highly productive habitats, competitively dominant species may monopolize space, excluding inferior competitors from the community. Biomass is high but species richness is low in such environments. Tropical rain forests are highly productive and have high biomass, they are relatively stable, and species richness is high. The majority of the plant family richness's examined in this study responded positively to high resource abundance indicated by high maximum NDVI values, suggesting that sites with potentially high productivity support more species than sites with lower potential. This is apparent as when the potentiality of productivity rises, the average variety of micronutrient combinations in fertile soils increases (Rosenweig and Abramsky 1993, Oindo and Skidmore 2002) leading to higher vegetative complexity (Bailey *et al.* 2004). The only negative response to maximum NDVI was obtained for the family Lentibulariaceae. This is ecologically reasonable, because the species of this family occur exclusively in shore habitats and shallow aquatic environments.

The shape of heterogeneity–richness relationships is generally positive (Gould 2000, Oindo and Skidmore 2002, Fairbanks and McGwire 2004, Levin *et al.* 2007). The variability of NDVI values usually reflects the heterogeneity of habitats, and is positively related with species richness of plants (Oindo and Skidmore 2002). However, in our study the responses between heterogeneity and species richness were mainly hump-shaped, indicating that the highest species richness occurs with moderate heterogeneity of productivity (Grime 1973, 1979). Our results showed that for example the family Salicaceae reaches its maximum species richness at intermediate levels of productivity and heterogeneity. The species of the Salicaceae family are well known to occur primarily in forests along riversides, and other places characterized by intermediate stages along the succession gradient in boreal forests. However, certain plant families in high latitudes appear to be specialized to the extremes of heterogeneity of primary productivity. For example the richness of the Orchidaceae family showed a clear nega-

tive relationship with heterogeneity. The species of the Orchidaceae family are rather demanding occurring strictly in stable old-growth forest, and a high level of heterogeneity may decrease habitat quality for them.

Conclusions

Developing conceptually simple models that are able to illustrate in a reliable manner how productivity and heterogeneity in productivity affect species richness may help the prediction of biodiversity in more remote and insufficiently surveyed areas. Such means can effectively decrease the need for extensive field sampling and can be essential for many conservation and management applications (Kerr and Ostrovsky 2003, Seto *et al.* 2004). Our approach, using continuous (unclassified) reflectance values provides a new possibility for researchers to develop relationships between species richness and remotely sensed data. Remotely sensed data allows trends over time to be examined, and areas of consistent high productive areas to be identified. Primary productivity and heterogeneity in the area, as estimated by using remotely sensed NDVI values, provide useful correlates of plant species richness in high-latitude areas. Spatial variability in species richness in 28 plant families was explained primarily by productivity, complemented by significant contributions from heterogeneity. Heterogeneity variables were often selected to species richness models at this scale and they showed mainly hump-shaped or linear responses.

These results have significant conservation implications. The role of both primary productivity and heterogeneity apparently was important in determining mesoscale species-richness patterns, more so than appears to be previously understood. Further work is needed to deepen our understanding of species density and to facilitate our ability to predict the responses of species richness to environmental changes. We conclude that remote-sensing based measures of productivity and heterogeneity have high potential as 'first filters' for identifying locations of high species richness in high-latitude forest landscapes.

Acknowledgements: Different parts of this research were funded by the Academy of Finland (project grant 116544) and EC FP6 Integrated Project ALARM (GOCE-CT-2003-506675). M. J. Bailey helped with correction of the English text. Two anonymous reviewers helped to improve a first draft of the manuscript. We thank P. Härmä for providing valuable remote sensing information.

References

- Airaksinen O. & Karttunen K. 2001. *Natura 2000 – habitats manual. Environment Guide 46*. Finnish Environment Institute, Helsinki.
- Atlas of Finland 1987. *Climate, Folio 131*. National Board of Survey & Geographical Society of Finland, Helsinki.
- Austin M.P. 2002. Spatial prediction of species distribution: an interface between ecological theory and statistical modelling. *Ecological Modelling* 157: 101–118.
- Bailey S.A., Horner-Devine M.C., Luck G., Moore L.A., Carney K.M., Anderson S., Betrus C. & Fleishman E. 2004. Primary productivity and species richness: relationships among functional guilds, residency groups and vagility classes at multiple spatial scales. *Ecography* 27: 207–217.
- Benayas J.M.R. & Scheiner S.M. 2002. Plant diversity, biogeography and environment in Iberia: Patterns and possible causal factors. *Journal of Vegetation Science* 13: 245–258.
- Bonan G.B. & Shugart H.H. 1989. Environmental factors and ecological processes in boreal forests. *Annual Review of Ecology and Systematics* 20: 1–28.
- Borcard D., Legendre P. & Drapeau P. 1992. Partialling out the spatial component of ecological variation. *Ecology* 73: 1045–1055.
- Box E.O., Holben B.N. & Kalb V. 1989. Accuracy of the AVHRR vegetation index as a predictor of biomass, primary productivity and net CO₂ flux. *Vegetatio* 80: 71–89.
- Bruun H.H. & Ejrnæs R. 2006. Community-level birth rate: a missing link between ecology, evolution and diversity. *Oikos* 113: 185–191.
- Currie D.J. 1991. Energy and large-scale patterns of animal- and plant-species richness. *American Naturalist* 137: 27–49.
- Currie D.J. & Paquin V. 1987. Large-scale biogeographical patterns of species richness of trees. *Nature* 329: 326–327.
- Dunning J.B., Danielson B.J. & Pulliam H.R. 1992. Ecological processes that effect populations in complex landscapes. *Oikos* 65: 169–175.
- Ekstrand S. 1996. Landsat TM-based forest damage assessment: correction for topographic effects. *Photogrammetric Engineering and Remote Sensing* 62: 151–161.
- Fairbanks D.H.K. & McGwire K.C. 2004. Patterns of floristic richness in vegetation communities of California: regional scale analysis with multi-temporal NDVI. *Global Ecology & Biogeography* 13: 221–235.
- Footy G.M. & Cutler M.E.J. 2003. Tree biodiversity in protected and logged Bornean tropical rain forests and

- its measurement by satellite remote sensing. *Journal of Biogeography* 30: 1053–1066.
- Gaston K.J. 2000. Global patterns in biodiversity. *Nature* 405: 220–227.
- Gillespie T.W. 2005. Predicting woody-plant species richness in tropical dry forests: a case study from south Florida, USA. *Ecological Applications* 15: 27–37.
- Gould W.A. 2000. Remote sensing of vegetation, plant species richness, and regional diversity hotspots. *Ecological Applications* 10: 1861–1870.
- Grace J.B. 1999. The factors controlling species density in herbaceous plant communities: an assessment. *Perspectives in Plant Ecology, Evolution and Systematics* 2: 1–28.
- Grime J.P. 1973. Competitive exclusion in herbaceous vegetation. *Nature* 242: 344–347.
- Grime J.P. 1979. *Plant strategies and vegetation processes*. John Wiley, Chichester.
- Grime J.P. 2001. *Plant strategies: vegetation processes and ecosystem properties*. John Wiley, Chichester.
- Guisan A., Edwards T.C.J. & Hastie T. 2002. Generalized linear and generalized additive models in studies of species distributions: setting the scene. *Ecological Modelling* 157: 89–100.
- H-Acevedo D. & Currie D.J. 2003. Does climate determine broad-scale patterns of species richness? A test of the causal link by natural experiment. *Global Ecology and Biogeography* 12: 461–473.
- Harrison S. & Grace J.B. 2007. Biogeographic affinity helps explain productivity–richness relationships at regional and local scales. *American Naturalist* 170: 5–15.
- Hastie T. & Tibshirani R. 1990. *Generalized additive models*. Chapman and Hall, London.
- Hawkins B.A. & Porter E.E. 2003. Does herbivore diversity depend on plant diversity? The case of Californian butterflies. *American Naturalist* 161: 40–49.
- Hawkins B.A., Field R., Cornell H.V., Currie D.J., Guégan J.-F., Kaufman D.M., Kerr J.T., Mittelbach G.G., Oberdorff T., O'Brien E.M., Porter E.E. & Turner J.R.G. 2003. Energy, water, and broad-scale geographic patterns of species richness. *Ecology* 84: 3105–3117.
- Heikkinen R.K. 1996. Predicting patterns of vascular plant species richness with composite variables: a meso-scale study in Finnish Lapland. *Vegetatio* 126: 151–165.
- Heikkinen R.K. 1998. Can richness patterns of rarities be predicted from mesoscale atlas data? A case study of vascular plants in the Kevo reserve. *Biological Conservation* 83: 133–143.
- Hill N.M. & Keddy P.A. 1992. Prediction of rarities from habitat variables: coastal plain plants on Nova Scotian lakeshores. *Ecology* 73: 1852–1859.
- Hjort J. & Luoto M. 2006. Modelling patterned ground distribution in Finnish Lapland: an integration of topographical, ground and remote sensing information. *Geografiska Annaler* 88A: 19–29.
- Huntley B., Berry P.M., Cramer W. & McDonald A.P. 1995. Modelling present and potential future ranges of some European higher plants using climate response surfaces. *Journal of Biogeography* 22: 967–1001.
- Hämet-Ahti L., Suominen J., Ulvinen T., Uotila P. & Vuokko S. 1986. *Retkeilykasvio*. Suomen Luonnonsuojelun Tuki Oy, Helsinki.
- Härmä P., Teiniranta R., Törmä M., Repo R., Järvenpää E. & Kallio M. 2004. Production of CORINE2000 land cover data using calibrated LANDSAT 7 ETM satellite image mosaics and digital Maps in Finland. In: *IEEE International Geoscience and Remote Sensing Symposium 20–24 September 2004, Anchorage, Alaska*, IEEE, pp. 2703–2706.
- Iverson L.R., Dale M.E., Scott C.T. & Prasad A. 1997. A GIS-derived integrated moisture index to predict forest composition and productivity of Ohio forests (U.S.A.). *Landscape Ecology* 12: 331–348.
- Kerr J. & Ostrovsky M. 2003. From space to species: ecological applications for remote sensing. *Trends in Ecology and Evolution* 18: 299–305.
- Kerr J.T. & Packer L. 1997. Habitat heterogeneity as a determinant of mammal species richness in high-energy regions. *Nature*: 252–254.
- Kerr J.T., Southwood T.R.E. & Cihlar J. 2001. Remotely sensed habitat diversity predicts butterfly species richness and community similarity in Canada. *Proceedings of the National Academy of Sciences USA* 98: 11365–11370.
- Kuusipalo J. 1984. Diversity pattern of the forest understorey vegetation in relation to some site characteristics. *Silva Fennica* 18: 121–131.
- Lehmann A., Overton J. & Leathwick J. 2002. GRASP: generalized regression analysis and spatial prediction. *Ecological Modelling* 157: 189–207.
- Levin N., Shmida A., Levanoni O., Tamari H. & Kark S. 2007. Predicting mountain plant richness and rarity from space using satellite-derived vegetation indices. *Diversity & Distributions* 13: 692–703.
- Lundholm J.T. & Larson D.W. 2003. Relationships between spatial environmental heterogeneity and plant species diversity on a limestone pavement. *Ecography* 26: 715–722.
- Luoto M. 2000. Modelling of rare plant species richness by landscape variables in an agriculture area in Finland. *Plant Ecology* 149: 157–168.
- Luoto M., Toivonen T. & Heikkinen R.K. 2002. Prediction of total and rare plant species richness in agricultural landscapes from satellite images and topographic data. *Landscape Ecology* 17: 195–217.
- McCullagh P. & Nelder J.A. 1989. *Generalized linear models*. Chapman & Hall, New York.
- Michalet R., Brooker R.W., Cavieres L.A., Kikvidze Z., Lortie C.J., Pugnaire F.I., Valiente-Banuet A. & Callaway R.M. 2006. Do biotic interactions shape both sides of the humped-back model of species richness in plant communities? *Ecology Letters* 9: 767–773.
- Mittelbach G.G., Steiner C.F., Scheiner S.M., Gross K.L., Reynolds H.L., Waide R.B., Willig M.R., Dodson S.I. & Gough L. 2001. What is the observed relationship between species richness and productivity? *Ecology* 82: 2381–2396.
- Muldavin E.H., Neville P. & Harper G. 2001. Indices of

- grassland biodiversity in the Chihuahuan desert ecoregion derived from remote sensing. *Conservation Biology* 15: 844–855.
- Nagendra H. 2001. Using remote sensing to assess biodiversity. *International Journal of Remote Sensing* 22: 2377–2400.
- Oindo B.O. & Skidmore A.K. 2002. Interannual variability of NDVI and species richness in Kenya. *International Journal of Remote Sensing* 23: 285–298.
- Opdam P. & Wascher D. 2004. Climate change meets habitat fragmentation: linking landscape and biogeographical scale levels in research and conservation. *Biological Conservation* 117: 285–297.
- Palmer M.W. 1994. Variation in species richness: towards a unification of hypotheses. *Folia Geobotanica et Phytotaxonomica* 29: 511–530.
- Palmer M.W., Earls P., Hoagland B.W., White P.S. & Wohlgenuth T. 2002. Quantitative tools for perfecting species lists. *Environmetrics* 13: 121–137.
- Parmesan C. 1996. Climate and species range. *Nature* 382: 765–766.
- Paruelo J.M., Epstein H.E., Lauenroth W.K. & Burke I.C. 1997. ANPP estimates from NDVI for the Central Grassland Region of the United States. *Ecology* 78: 953–958.
- Parviainen M., Luoto M., Rytteri T. & Heikkinen R.K. 2008. Modelling the occurrence of threatened plant species in taiga landscapes: methodological and ecological perspectives. *Journal of Biogeography* 35: 1888–1905.
- Pearson R.G., Dawson T.P. & Liu C. 2004. Modelling species distributions in Britain: a hierarchical integration of climate and land-cover data. *Ecography* 27: 285–298.
- Pettorelli N., Vik J.O., Mysterud A., Gaillard J.-M., Tucker C.J. & Stenseth N.C. 2005. Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution* 20: 503–510.
- Pärtel M., Laanisto L. & Zobel M. 2007. Contrasting plant productivity–diversity relationships across latitude: the role of evolutionary history. *Ecology* 88: 1091–1097.
- Rahbek C. & Graves G.R. 2001. Multiscale assessment of patterns of avian species richness. *Proceedings of the National Academy of Sciences USA* 98: 4534–4539.
- Reed B.C., Brown J.F., Vander Zee D., Loveland T.R., Merchant J.W. & Ohlen D.O. 1994. Measuring Phenological Variability from Satellite Imagery. *Journal of Vegetation Science* 5: 703–714.
- Ricklefs R.E. 1977. Environmental heterogeneity and plant species diversity: a hypothesis. *American Naturalist* 111: 376–381.
- Rocchini D. 2007. Effects of spatial and spectral resolution in estimating ecosystem α -diversity by satellite imagery. *Remote Sensing of Environment* 111: 423–434.
- Rocchini D., Chiarucci A. & Loiselle S.A. 2004. Testing the spectral variation hypothesis by using satellite multi-spectral images. *Acta Oecologica* 26: 117–120.
- Rocchini D., Andreini Butini S. & Chiarucci A. 2005. Maximizing plant species inventory efficiency by means of remotely sensed spectral distances. *Global Ecology and Biogeography* 14: 431–437.
- Rosenweig M.L. 1995. *Species diversity in space and time*. Cambridge University Press, Cambridge.
- Rosenweig M.L. & Abramsky Z. (eds.) 1993. *How are diversity and productivity related? Species diversity in ecological communities*. University of Chicago Press, Chicago.
- Rosenzweig M.L. 1995. *Species diversity in space and time*. Cambridge University Press, Cambridge.
- Rouse J.W.Jr., Hass R.H., Deering D.W., Schell J.A. & Harlan J.C. 1973. *Monitoring the vernal advancement and retrogression (green wave effect) of natural vegetation*. NASA/GSFC Type III Final report. Geenbelt, Maryland, MD.
- Seto K.C., Fleishman E., Fay J.P. & Betrus C.J. 2004. Linking spatial patterns of bird and butterfly species richness with Landsat TM derived NDVI. *International Journal of Remote Sensing* 25: 4309–4324.
- Stoms D.M. 1994. Scale dependence of species richness maps. *Professional Geographer* 46: 346–358.
- Söyrinki N. & Saari V. 1980. Die Flora im Oulanka Nationalpark, Nordfinland. *Acta Florestica Fennica* 114: 1–149.
- Thuiller W., Araújo M.B. & Lavorel S. 2004. Do we need land-cover data to predict species distributions in Europe? *Journal of Biogeography* 31: 353–361.
- Tilman D. 1982. *Resource competition and community structure*. Princeton University Press, Princeton.
- Tucker C.J. 1979. Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment* 8: 127–150.
- Tucker C.J. & Sellers P.J. 1986. Satellite remote-sensing of primary production. *International Journal of Remote Sensing* 7: 1395–1416.
- Waide R.B., Willig M.R., Steiner C.F., Mittelbach G., Gough L., Dodson S.I., Juday G.P. & Parmenter R. 1999. The relationship between productivity and species richness. *Annual Review of Ecological Systems* 30: 257–300.
- VanderMeulen M.A., Hudson A.J. & Scheiner S.M. 2001. Three evolutionary hypotheses for the hump-shaped productivity/diversity curve. *Evolutionary Ecology Research* 3: 379–392.
- Vasari Y., Tonkov S., Vasari A. & Nikolova A. 1996. The Late-quaternary history of the vegetation and flora in northeastern Finland in the light of a re-investigation of Aapalampi in Salla. *Auqilo Series Botany* 36: 27–41.
- Venables W.N. & Ripley B.D. 2002. *Modern applied statistics with S*. Springer-Verlag, Berlin.
- Whittaker R.J. & Heegaard E. 2003. What is the observed relationship between species richness and productivity? Comment. *Ecology* 84: 3384–3390.
- Wiens J.J. & Donoghue M.J. 2004. Historical biogeography, ecology and species richness. *Trends in Ecology & Evolution* 19: 639–644.
- Virtanen R., Oksanen J., Oksanen L. & Razzhivin V.Y. 2006. Broad-scale vegetation–environment relationships in Eurasian high-latitude areas. *Journal of Vegetation Science* 17: 519–528.
- Wood S. & Augustin N. 2002. GAMs with integrated model selection using penalized regression splines and applications to environmental modelling. *Ecological Modelling* 157: 157–177.
- Wright D.H., Currie D.J. & Maurer B.A. 1993. Energy

supply and patterns of species richness on local and regional scales. In: Ricklefs R.E. & Schluter D. (eds.), *Species diversity in ecological communities: historical and geographical perspectives*, University of Chicago

Press, pp. 66–74.

Yee T.W. & Mitchell N.D. 1991. Generalized additive models in plant ecology. *Journal of Vegetation Science* 2: 587–602.

Appendix 1. A complete list of vascular plant species.

Species	Number of occupied cells	Species	Number of occupied cells
Lycopodiaceae		<i>Ranunculus peltatus</i>	17
<i>Huperzia selago</i>	107	<i>Ranunculus trichophyllus</i> ssp. <i>trichophyllus</i>	2
<i>Lycopodiella inundata</i>	1	<i>Ranunculus trichophyllus</i> ssp. <i>eradicatus</i>	10
<i>Lycopodium annotinum</i>	382	<i>Thalictrum flavum</i>	102
<i>Lycopodium clavatum</i>	19	Betulaceae	
<i>Diphasium complanatum</i>	332	<i>Betula pendula</i>	396
Equisetaceae		<i>Betula pubescens</i>	438
<i>Equisetum hyemale</i>	113	<i>Betula nana</i>	330
<i>Equisetum variegatum</i>	76	<i>Alnus incana</i>	350
<i>Equisetum scirpoides</i>	248	Caryophyllaceae	
<i>Equisetum fluviatile</i>	298	<i>Arenaria ciliata</i> ssp. <i>pseudofrigida</i>	13
<i>Equisetum palustre</i>	263	<i>Stellaria media</i>	2
<i>Equisetum sylvaticum</i>	318	<i>Stellaria graminea</i>	63
<i>Equisetum pretense</i>	297	<i>Stellaria longifolia</i>	9
<i>Equisetum arvense</i>	208	<i>Stellaria calycantha</i>	45
Dryopteridaceae		<i>Stellaria crassifolia</i>	15
<i>Athyrium filix-femina</i>	43	<i>Cerastium alpinum</i>	45
<i>Diplazium sibiricum</i>	18	<i>Cerastium fontanum</i> ssp. <i>scandicum</i>	78
<i>Cystopteris fragilis</i> var. <i>fragilis</i>	42	<i>Cerastium fontanum</i> ssp. <i>vulgare</i>	30
<i>Cystopteris fragilis</i> var. <i>dickieana</i>	48	<i>Sagina procumbens</i>	3
<i>Cystopteris montana</i>	90	<i>Scleranthus annuus</i>	1
<i>Woodsia ilvensis</i>	20	<i>Spergula arvensis</i>	1
<i>Woodsia alpina</i>	46	<i>Lychnis alpina</i>	1
<i>Woodsia glabella</i>	55	<i>Silene tatarica</i>	10
<i>Matteuccia struthiopteris</i>	37	<i>Silene vulgaris</i>	1
<i>Dryopteris carthusiana</i>	11	<i>Silene latifolia</i> ssp. <i>alba</i>	1
<i>Gymnocarpium dryopteris</i>	335	<i>Silene dioica</i>	1
<i>Gymnocarpium continetale</i>	40	<i>Gypsophila fastigiata</i>	15
<i>Gymnocarpium robertianum</i>	37	<i>Dianthus superbus</i>	53
Ranunculaceae		Polygonaceae	
<i>Trollius europaeus</i>	251	<i>Polygonum aviculare</i>	1
<i>Actaea erythrocarpa</i>	155	<i>Polygonum lapathifolium</i>	1
<i>Caltha palustris</i>	191	<i>Polygonum amphibium</i>	5
<i>Ranunculus repens</i>	125	<i>Polygonum viviparum</i>	246
<i>Ranunculus acris</i>	160	<i>Fallopia convolvulus</i>	1
<i>Ranunculus monophyllus</i>	10	<i>Rumex acetosella</i>	32
<i>Ranunculus auricomus</i>	107	<i>Rumex acetosa</i>	11
<i>Ranunculus hyperboreus</i>	2	<i>Rumex aquaticus</i>	9
<i>Ranunculus lapponicus</i>	2	<i>Rumex longifolius</i>	5
<i>Ranunculus reptans</i>	104		

continued

Appendix 1. Continued.

Species	Number of occupied cells	Species	Number of occupied cells
Violaceae			
<i>Viola rupestris</i>	70	<i>Pyrola rotundifolia</i>	190
<i>Viola riviniana</i>	8	<i>Orthilia secunda</i>	365
<i>Viola canina</i> ssp. <i>montana</i>	154	<i>Moneses uniflora</i>	237
<i>Viola epipsila</i>	313	Saxifragaceae	
<i>Viola selkirkii</i>	36	<i>Saxifraga nivalis</i>	68
Brassicaceae			
<i>Sisymbrium altissimum</i>	1	<i>Saxifraga hirculus</i>	19
<i>Erysimum hieraciifolium</i>	28	<i>Saxifraga aizoides</i>	9
<i>Erysimum cheiranthoides</i> ssp. <i>cheiranthoides</i>	1	<i>Saxifraga cespitosa</i>	6
<i>Erysimum cheiranthoides</i> ssp. <i>altum</i>	1	Rosaceae	
<i>Barbarea stricta</i>	15	<i>Filipendula ulmaria</i>	322
<i>Rorippa palustris</i>	10	<i>Rubus chamaemorus</i>	351
<i>Cardamine pratensis</i> ssp. <i>dentata</i>	136	<i>Rubus arcticus</i>	112
<i>Draba norvegica</i>	1	<i>Rubus saxatilis</i>	359
<i>Draba cinerea</i>	20	<i>Rubus idaeus</i>	145
<i>Capsella bursa-pastoris</i>	2	<i>Rosa majalis</i>	141
<i>Thlaspi arvense</i>	1	<i>Dryas octopetala</i>	17
<i>Subularia aquatica</i>	25	<i>Geum rivale</i>	299
Salicaceae			
<i>Salix pentandra</i>	25	<i>Potentilla palustris</i>	314
<i>Salix reticulata</i>	3	<i>Potentilla nivea</i>	13
<i>Salix myrsinites</i>	186	<i>Potentilla norvegica</i>	2
<i>Salix glauca</i>	75	<i>Potentilla erecta</i>	164
<i>Salix phylicifolia</i>	375	<i>Fragaria vesca</i>	75
<i>Salix myrsinifolia</i>	152	<i>Alchemilla glomerulans</i>	1
<i>Salix aurita</i>	5	<i>Sorbus aaucuparia</i>	377
<i>Salix caprea</i>	408	<i>Prunus padus</i>	191
<i>Salix starkeana</i>	22	Fabaceae	
<i>Salix xerophila</i>	8	<i>Astragalus frigidus</i>	79
<i>Salix myrtilloides</i>	204	<i>Astragalus alpinus</i>	74
<i>Salix repens</i>	1	<i>Oxytropis campestris</i>	49
<i>Salix lapponum</i>	292	<i>Vicia cracca</i>	3
<i>Salix hastata</i>	190	<i>Vicia sylvatica</i>	1
<i>Salix pyrolifolia</i>	1	<i>Trifolium repens</i>	23
<i>Populus tremula</i>	343	<i>Trifolium hybridum</i>	2
Ericaceae			
<i>Calluna vulgaris</i>	421	<i>Trifolium pratense</i>	11
<i>Rhododendron tomentosum</i>	438	<i>Anthyllis vulneraria</i>	3
<i>Arctostaphylos uva-ursi</i>	130	Onagraceae	
<i>Arctostaphylos alpina</i>	2	<i>Cincaea alpina</i>	9
<i>Andromeda polifolia</i>	322	<i>Epilobium angustifolium</i>	360
<i>Vaccinium oxycoccus</i>	279	<i>Epilobium palustre</i>	154
<i>Vaccinium microcarpum</i>	295	<i>Epilobium davuricum</i>	72
<i>Vaccinium vitis-idaea</i>	440	<i>Epilobium hornemannii</i>	29
<i>Vaccinium uliginosum</i>	439	<i>Epilobium alsinifolium</i>	16
<i>Vaccinium myrtilus</i>	440	Apiaceae	
Pyrolaceae			
<i>Pyrola minor</i>	292	<i>Anthriscus sylvestris</i>	56
<i>Pyrola media</i>	12	<i>Cicuta vivosa</i>	2
<i>Pyrola chlorantha</i>	57	<i>Carum carvi</i>	2
		<i>Angelica sylvestris</i>	245
		<i>Peucedanum palustre</i>	19
		<i>Heracleum sphondylium</i> ssp. <i>sibiricum</i>	1

continued

Appendix 1. Continued.

Species	Number of occupied cells	Species	Number of occupied cells
Rubiaceae			
<i>Galium boreale</i>	168	<i>Saussurea alpina</i>	326
<i>Galium triflorum</i>	2	<i>Cirsium helenioides</i>	301
<i>Galium uliginosum</i>	268	<i>Cirsium palustre</i>	1
<i>Galium palustre</i>	184	Cichoriaceae	
<i>Galium trifidum</i>	14	<i>Leontodon autumnalis</i>	8
Lamiaceae			
<i>Scutellaria galericulata</i>	100	<i>Sonchus asper</i>	1
<i>Galeopsis speciosa</i>	1	<i>Lactuca sibirica</i>	4
<i>Galeopsis bifida</i>	4	<i>Cicerbita alpina</i>	12
<i>Prunella vulgaris</i>	1	<i>Taraxatum</i> ssp.	189
<i>Thymus sepyllum</i> ssp. <i>tanaënsis</i>	70	<i>Lapsana communis</i>	1
Scrophulariaceae			
<i>Veronica serpyllifolia</i> ssp. <i>serpyllifolia</i>	6	<i>Crepis paludosa</i>	219
<i>Veronica scutellata</i>	17	<i>Crepis tectorum</i> ssp. <i>tectorum</i>	3
<i>Veronica longifolia</i>	106	<i>Crepis tectorum</i> ssp. <i>nigrescens</i>	2
<i>Melampyrum sylvaticum</i>	309	<i>Hieracium</i> ssp.	349
<i>Melampyrum pratense</i>	430	<i>Hieracium umbellatum</i>	45
<i>Euphrasia frigida</i>	140	Orchidaceae	
<i>Bartsia alpina</i>	211	<i>Cypripedium calceolus</i>	59
<i>Pedicularis sceptrum-carolinum</i>	50	<i>Epipactis atrorubens</i>	30
<i>Pedicularis palustris</i> ssp. <i>borealis</i>	203	<i>Epipogium aphyllum</i>	4
<i>Rhinanthus minor</i>	47	<i>Listera ovata</i>	13
<i>Rhinanthus serotinus</i>	1	<i>Listera cordata</i>	118
Lentibulariaceae			
<i>Pinguicula alpina</i>	132	<i>Goodyera repens</i>	75
<i>Pinguicula villosa</i>	34	<i>Corallorhiza trifida</i>	52
<i>Pinguicula vulgaris</i>	118	<i>Calypso bulbosa</i>	56
<i>Utricularia minor</i>	28	<i>Coeloglossum viride</i>	126
<i>Utricularia intermedia</i>	125	<i>Gymnadenia conopsea</i>	162
<i>Utricularia vulgaris</i>	27	<i>Dactylorhiza incarnate</i>	41
Asteraceae			
<i>Solidago virgaurea</i>	424	<i>Dactylorhiza traunsteineri</i>	35
<i>Erigeron acer</i> ssp. <i>acer</i>	3	<i>Dactylorhiza maculata</i>	193
<i>Erigeron acer</i> ssp. <i>politus</i>	13	Potamogetonaceae	
<i>Erigeron acer</i> ssp. <i>decoloratus</i>	34	<i>Potamogeton filiformis</i>	8
<i>Gnaphalium sylvaticum</i>	7	<i>Potamogeton compressus</i>	1
<i>Antennaria dioica</i>	195	<i>Potamogeton berchtoldii</i>	19
<i>Inula salicina</i>	10	<i>Potamogeton alpinus</i>	87
<i>Achillea ptarmica</i>	2	<i>Potamogeton natans</i>	31
<i>Achillea millefolium</i>	68	<i>Potamogeton gramineus</i>	84
<i>Tripleurospermum maritimum</i>	3	<i>Potamogeton lucens</i>	2
<i>Tripleurospermum inodorum</i>	3	<i>Potamogeton praelongus</i>	2
<i>Matricaria matricarioides</i>	2	<i>Potamogeton perfoliatus</i>	46
<i>Tanacetum vulgare</i>	2	Juncaceae	
<i>Artemisia vulgaris</i>	1	<i>Juncus bufonius</i>	2
<i>Tussilago farfara</i>	13	<i>Juncus alpinus</i> ssp. <i>nodulosus</i>	43
<i>Petasites frigidus</i>	83	<i>Juncus filiformis</i>	105
<i>Arnica angustifolia</i> ssp. <i>alpina</i>	10	<i>Juncus stygius</i>	111
<i>Senecio vernalis</i>	1	<i>Juncus triglumis</i>	28
<i>Senecio vulgaris</i>	1	<i>Luzula pilosa</i>	412
		<i>Luzula multiflora</i> ssp. <i>multiflora</i>	3
		<i>Luzula multiflora</i> ssp. <i>frigida</i>	36
		<i>Luzula sudentica</i>	66
		<i>Luzula pallescens</i>	31

continued

Appendix 1. Continued.

Species	Number of occupied cells	Species	Number of occupied cells
Cyperaceae		<i>Carex serotina</i> ssp. <i>pulchella</i>	1
<i>Eriophorum vaginatum</i>	320	<i>Carex lasiocarpa</i>	274
<i>Eriophorum brachyantherum</i>	25	<i>Carex vesicaria</i>	103
<i>Eriophorum russeolum</i>	3	<i>Carex rotundata</i>	2
<i>Eriophorum scheuchzeri</i>	3	<i>Carex rostrata</i>	325
<i>Eriophorum angustifolium</i>	260	Poaceae	
<i>Eriophorum latifolium</i>	182	<i>Molinia caerulea</i>	334
<i>Eriophorum gracile</i>	13	<i>Phragmites australis</i>	119
<i>Schoenoplectus lacustris</i>	14	<i>Nardus stricta</i>	2
<i>Trichophorum alpinum</i>	249	<i>Melica nutans</i>	310
<i>Trichophorum cespitosum</i>	259	<i>Festuca ovina</i>	278
<i>Eleocharis palustris</i> ssp. <i>palustris</i>	1	<i>Festuca rubra</i>	143
<i>Eleocharis acicularis</i>	10	<i>Festuca pratensis</i>	2
<i>Eleocharis quinqueflora</i>	23	<i>Lolium perenne</i>	2
<i>Schoenus ferrugineus</i>	9	<i>Lolium multiflorum</i>	1
<i>Rhynchospora alba</i>	1	<i>Poa annua</i>	13
<i>Carex capitata</i>	92	<i>Poa trivialis</i>	12
<i>Carex pauciflora</i>	195	<i>Poa alpina</i>	38
<i>Carex chordorrhiza</i>	248	<i>Poa pratensis</i>	1
<i>Carex dioica</i>	282	<i>Poa subcaerulea</i>	3
<i>Carex heleonastes</i>	18	<i>Poa alpigena</i>	211
<i>Carex canescens</i>	283	<i>Poa nemoralis</i>	145
<i>Carex brunnescens</i>	109	<i>Poa glauca</i>	9
<i>Carex disperma</i>	2	<i>Poa palustris</i>	5
<i>Carex tenuiflora</i>	10	<i>Dactylis glomerata</i>	1
<i>Carex loliacea</i>	131	<i>Deschampsia cespitosa</i>	268
<i>Carex elongata</i>	1	<i>Deschampsia flexuosa</i>	437
<i>Carex echinata</i>	15	<i>Calamagrostis epigejos</i>	210
<i>Carex diandra</i>	46	<i>Calamagrostis canescens</i>	5
<i>Carex appropinquata</i>	15	<i>Calamagrostis purpurea</i> ssp. <i>phragmitoides</i>	364
<i>Carex aquatilis</i>	28	<i>Calamagrostis stricta</i>	49
<i>Carex acuta</i>	109	<i>Calamagrostis lapponica</i>	309
<i>Carex nigra</i> ssp. <i>nigra</i>	4	<i>Agrostis stolonifera</i>	29
<i>Carex nigra</i> ssp. <i>juncella</i>	295	<i>Agrostis gigantea</i>	8
<i>Carex cespitosa</i>	244	<i>Agrostis capillaris</i>	134
<i>Carex elata</i> ssp. <i>omskiana</i>	1	<i>Agrostis canina</i>	3
<i>Carex buxbaumii</i> ssp. <i>buxbaumii</i>	128	<i>Agrostis mertensii</i>	193
<i>Carex buxbaumii</i> ssp. <i>mutica</i>	7	<i>Alopecurus pratensis</i>	4
<i>Carex norvegica</i> ssp. <i>inferalpina</i>	112	<i>Alopecurus arundinaceus</i>	1
<i>Carex vaginata</i>	259	<i>Alopecurus geniculatus</i>	1
<i>Carex panicea</i>	91	<i>Alopecurus aequalis</i>	20
<i>Carex livida</i>	82	<i>Phleum pratense</i>	8
<i>Carex magellanica</i> ssp. <i>irrigua</i>	288	<i>Phleum alpinum</i>	136
<i>Carex limosa</i>	223	<i>Phalaris arundinacea</i>	187
<i>Carex rariflora</i>	1	<i>Hierochloë hirta</i> ssp. <i>arctica</i>	68
<i>Carex laxa</i>	17	<i>Anthoxanthum odoratum</i>	136
<i>Carex globularis</i>	389	<i>Milium effusum</i>	104
<i>Carex ericetorum</i>	45	<i>Bromus hordeaceus</i>	2
<i>Carex digitata</i>	76	<i>Elymus repens</i>	1
<i>Carex glacialis</i>	10	<i>Elymus caninus</i>	195
<i>Carex capillaris</i>	165	<i>Elymus fibrosus</i>	8
<i>Carex flava</i>	228	<i>Elymys alaskanus</i>	45
<i>Carex jemtlandica</i>	14	<i>Hordeum jubatum</i>	1
<i>Carex bergrothii</i>	27		

Appendix 2. Explained deviance and cross-validated (cvCOR) values of the productivity, heterogeneity and hybrid (productivity + heterogeneity) model. cvCOR values were calculated as the Pearson correlation index between the observed and predicted values of species richness.

Family	Productivity model		Heterogeneity model		Hybrid model	
	Explained deviance (%)	cvCOR	Explained deviance (%)	cvCOR	Explained deviance (%)	cvCOR
Lycopodiaceae	12.824	0.372	3.756	0.177	12.824	0.372
Equisetaceae	7.086	0.254	9.886	0.298	13.346	0.366
Dryopteridaceae	29.012	0.505	11.860	0.307	31.367	0.502
Ranunculaceae	37.715	0.632	18.962	0.439	39.993	0.644
Betulaceae	9.663	0.278			9.663	0.278
Caryophyllaceae	33.142	0.525	17.896	0.347	35.656	0.536
Polygonaceae	15.657	0.334	8.735	0.284	16.706	0.398
Violaceae	19.064	0.484	14.267	0.401	20.869	0.506
Brassicaceae	39.970	0.552	24.749	0.404	42.128	0.576
Salicaceae	10.656	0.294	8.725	0.255	16.804	0.337
Ericaceae	7.567	0.268	7.416	0.206	15.532	0.373
Pyrolaceae	17.855	0.453	7.923	0.272	18.643	0.45
Saxifragaceae	20.253	0.322	12.853	0.26	23.592	0.382
Rosaceae	21.006	0.462	12.527	0.34	24.992	0.491
Fabaceae	32.348	0.527	16.331	0.304	34.516	0.548
Onagraceae	18.186	0.431	10.213	0.313	20.808	0.463
Apiaceae	12.960	0.404	8.748	0.347	15.303	0.447
Rubiaceae	16.366	0.452	11.778	0.374	20.814	0.484
Lamiaceae	38.126	0.586	23.784	0.444	41.050	0.603
Scrophulariaceae	21.014	0.46	18.015	0.421	25.185	0.504
Lentibulariaceae	1.970	0.0853	7.512	0.209	8.007	0.264
Asteraceae	23.842	0.487	10.049	0.299	24.912	0.494
Cichoriaceae	23.689	0.492	9.656	0.336	25.068	0.507
Orchidaceae	5.521	0.18	3.151	0.0261	7.142	0.18
Potamogetonaceae	26.517	0.471	30.635	0.524	36.159	0.569
Juncaceae	10.708	0.305	12.119	0.318	15.436	0.372
Cyperaceae	2.716	0.0326	8.952	0.269	14.020	0.316
Poaceae	29.346	0.539	18.625	0.419	33.553	0.57
Mean	19.456	0.400	12.930	0.318	23.003	0.448
SD	10.847	0.147	6.433	0.099	10.291	0.112